

|  |  |                                    |  |  |
|--|--|------------------------------------|--|--|
| <b>1. AGENCY USE ONLY (Leave blank)</b>  |  | <b>2. REPORT DATE</b><br>June 2000 | <b>3. REPORT TYPE AND DATES COVERED</b><br>Final (15 May 97 - 14 May 00) |  |
| <b>4. TITLE AND SUBTITLE</b><br>Function of Protein Phosphatase 2A in Control of Proliferation: Isolation and Analysis of Dominant-Defective Mutants             |  |                                    | <b>5. FUNDING NUMBERS</b><br>DAMD17-97-1-7113                            |  |
| <b>6. AUTHOR(S)</b><br>Alison DeLong, Ph.D.  |  |                                    |  |  |
| <b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b><br>Brown University<br>Providence, Rhode Island 02912<br><br><b>E-MAIL:</b><br>Alison_DeLong@Brown.Edu |  |                                    | <b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>                          |  |
| <b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b><br><br>U.S. Army Medical Research and Materiel Command<br>Fort Detrick, Maryland 21702-5012     |  |                                    | <b>10. SPONSORING / MONITORING AGENCY REPORT NUMBER</b>                  |  |
| <b>11. SUPPLEMENTARY NOTES</b>   |  |                                    |  |  |
| <b>12a. DISTRIBUTION / AVAILABILITY STATEMENT</b><br>Approved for public release; distribution unlimited   |  |                                    | <b>12b. DISTRIBUTION CODE</b><br><br><b>20010228 087</b>                 |  |
| <b>13. ABSTRACT (Maximum 200 Words)</b>  |  |                                    |  |  |

Reversible protein phosphorylation plays a crucial role in the circuitry controlling diverse cellular processes, and the activities of phosphorylating and dephosphorylating enzymes must be carefully balanced in normal cells. It is well documented that many of these enzymes are constitutively expressed, but their activities are tightly regulated by a variety of post translational mechanisms. In the case of the serine/threonine-specific protein phosphatase 2A (PP2A), a catalytic subunit is bound by two regulatory subunits designated A and B. Previously, we established a functional complementation assay for PP2A-C in the yeast *S. cerevisiae* and used this system to isolate two dominant-defective mutants in the human PP2A-C $\alpha$  gene. We generated epitope-tagged forms of the wild-type and mutant alleles and showed that the epitope-tagged alleles retain biological function in our yeast system. The mutant alleles show reduced binding of the A regulatory subunit in yeast and mammalian cells. Stable fibroblastic cell lines expressing mutant and wild-type proteins have been isolated. These studies will thus increase our understanding of PP2A function in the regulation of proliferation and malignant transformation, and will yield important information about structure-function relationships in the PP2A catalytic and regulatory subunit proteins.

|  |   |  |  |
|--|---|--|--|
| <b>14. SUBJECT TERMS</b><br>Breast Cancer, Idea Award Protein Phosphatase 2A / Functional Complementation in <i>S. cerevisiae</i> / Dominant-Defective and Gain-of-function Mutants / Growth Regulation / Cell Culture |   |  | <b>15. NUMBER OF PAGES</b><br>21               |
|  |   |  | <b>16. PRICE CODE</b>                          |
| <b>17. SECURITY CLASSIFICATION OF REPORT</b><br>Unclassified   | <b>18. SECURITY CLASSIFICATION OF THIS PAGE</b><br>Unclassified | <b>19. SECURITY CLASSIFICATION OF ABSTRACT</b><br>Unclassified | <b>20. LIMITATION OF ABSTRACT</b><br>Unlimited |

AD \_\_\_\_\_

Award Number: DAMD17-97-1-7113

TITLE: Function of Protein Phosphatase 2A in Control of  
Proliferation: Isolation and Analysis of Dominant-  
Defective Mutants

PRINCIPAL INVESTIGATOR: Alison DeLong, Ph.D.

CONTRACTING ORGANIZATION: Brown University  
Providence, Rhode Island 02912

REPORT DATE: June 2000

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

## FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

\_\_\_ Where copyrighted material is quoted, permission has been obtained to use such material.

\_\_\_ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

\_\_\_ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

N/A In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

\_\_\_ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

X In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

X In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N/A In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Lisa D. Brown 6/12/00

## TABLE OF CONTENTS

|                                   |   |
|-----------------------------------|---|
| Front cover.....                  | 1 |
| Standard Form 298.....            | 2 |
| Foreword.....                     | 3 |
| Table of Contents.....            | 4 |
| Introduction.....                 | 5 |
| Body.....                         | 5 |
| Key Research Accomplishments..... | 6 |
| Reportable Outcomes.....          | 7 |
| Conclusions.....                  | 7 |
| References.....                   | 8 |
| Appendices.....                   | 9 |

## INTRODUCTION

Protein phosphatase 2A (PP2A) comprises catalytic (C) and regulatory (A and B) subunits, and a heterotrimeric (ABC) holoenzyme is thought to predominate *in vivo*, although a heterodimeric (AC) form also has been purified (reviewed in Walter and Mumby, 1993). Although the total number of PP2A substrates is not known, it is likely that regulation of PP2A activity is necessary for the correct functioning of signal transduction cascades in which PP2A participates. Essential PP2A subunit functions are strongly conserved throughout evolution, as evidenced by the ability of an *Arabidopsis* PP2A-A subunit gene (*RCN1*) to complement the temperature-sensitive phenotype of a null mutation in the yeast PP2A-A subunit gene (*tpd3*; Garbers et al., 1996). PP2A has not been identified as an oncogene per se, but its interactions with proteins responsible for tumorigenesis are of demonstrated functional significance. Evidence for the role of PP2A in mammalian cell proliferation has originated in studies showing that PP2A is a target for the small-T antigen of SV40 and the small- and medium-T antigens of polyoma virus. These tumor antigens bind the AC heterodimeric PP2A complex (Pallas et al., 1990; Walter et al., 1990) and exclude the B subunit. Interaction with SV40 small t antigen reduces PP2A activity toward the mitogen-activated protein kinases (MAPKs) ERK1 and MEK1, causing deregulation of the MAPK cascade and induction of cell growth (Sontag et al., 1993). Middle T antigen mutants that fail to bind PP2A also are defective in transformation (Campbell et al., 1995). Furthermore, the tumor-promoting agent okadaic acid (OKA; Suganuma et al., 1988) is a potent and specific inhibitor of PP2A and PP1, with PP2A showing approximately 100-fold higher sensitivity (Cohen et al., 1989). The targeting of PP2A by viral T antigens and the tumor-promoting effect of OKA at low concentrations both support the hypothesis that PP2A activity normally suppresses proliferation, and that antagonism of PP2A may be important in tumorigenesis.

Dominant gain-of-function alleles have revealed the identities and functions of many proteins involved in growth control. We proposed to isolate dominant-defective mutants of human PP2A as a tool for investigating potential growth control functions of PP2A in normal and breast cancer cells. Possible causes of dominance for such mutants include competition with the wild-type protein for substrates or for positive regulatory subunits. The PP2A catalytic subunit (PP2A-C) provides a good target for mutagenesis aimed at generation of dominant alleles, because putative catalytic residues have been identified and interactions with positive regulatory subunits are known to be important for normal activity.

Previously, we established a functional complementation assay for PP2A-C in the yeast *S. cerevisiae* and used this system to isolate two dominant-defective mutants in the human PP2A-C $\alpha$  gene. We had isolated epitope-tagged forms of the wild-type and mutant alleles and tested those alleles for biological function in our yeast system. Our recent work has been focused on isolation and characterization of stably transformed cell lines expressing epitope-tagged mutant and wild-type PP2A-C alleles, and analysis of PP2A subunit interactions in yeast and mammalian cells via immunoprecipitation.

## BODY

### *Analysis of the dominant defective phenotypes of PP2A-C $\alpha$ mutants H118N and R89A*

We have shown that *S. cerevisiae* can be used as a rapid *in vivo* assay system for human PP2A-C activity. Mutation of a conserved histidine residue in the C $\alpha$  sequence (the C $\alpha$ -H118N mutant) confers a strong dominant-defective phenotype; mutation of a conserved

arginine residue (the C $\alpha$ -R89A mutant) results in a weaker dominant-defective phenotype. A detailed description of this work was published last year in *Gene*; a reprint is appended.

We have considered several alternative hypotheses that might explain the dominance of the H118N and R89A mutants in yeast cells. First the mutant enzyme might bind substrate efficiently and prevent the Pph21-ts gene product from interacting with one or more essential substrate(s). Second, the mutant enzyme might interact normally with PP2A regulatory subunits (A and/or B) that are required for normal activity of the Pph21 gene product. Third, the mutant enzyme might bind another protein (neither substrate nor classical regulatory subunit) not bound by the wild-type enzyme. We have already demonstrated that expression of the heterologous PP2A-C does not down-regulate the endogenous PP2A-C expression.

We have used immunoprecipitation to assay binding of the A regulatory subunit to mutant and wild-type C $\alpha$  subunits in yeast and mammalian cells. Anti-FLAG antibody efficiently precipitates the epitope-tagged proteins from cells expressing FLAG-C $\alpha$ <sup>+</sup>, FLAG-C $\alpha$ -H118N, FLAG-C $\alpha$  R89A and FLAG-C $\alpha$ -R89A/H118N proteins. However, the endogenous yeast A regulatory subunit is co-immunoprecipitated efficiently only in extracts from cells expressing the wild-type FLAG-C $\alpha$ <sup>+</sup> protein. The H118N, R89A and double mutant R89A/H118N mutations markedly reduce the amount of co-immunoprecipitating A subunit, suggesting that these lesions weaken the association of the C $\alpha$  subunit with the A regulatory subunit. To assay for a similar effect in mammalian cells, COS cells transfected with FLAG-C $\alpha$ <sup>+</sup> and FLAG-C $\alpha$ -H118N expression constructs were also tested for co-immunoprecipitation of endogenous A subunit using the anti-FLAG antibody. While the A subunit was easily detected in immunoprecipitates from cells expressing the wild-type C $\alpha$ <sup>+</sup> subunit, A subunit co-immunoprecipitation was markedly reduced in extracts from cells expressing C $\alpha$ -H118N. Because A subunit binding is required for formation of heterotrimeric PP2A complexes (i.e. subsequent B subunit binding; Ruediger et al., 1992), these results suggest that the protein products of the mutant alleles do not compete with endogenous C subunits for A and B subunit binding. Thus the dominant-defective phenotype is not due to competition for regulatory subunit binding. We are currently generating the R89A expression constructs required to test for co-immunoprecipitation of mammalian A subunit with the R89A mutant in COS cells. We have attempted to assay for co-immunoprecipitation of the yeast Tap42 protein (homolog of the mammalian  $\alpha$ 4 protein; Di Como et al. 1996), but due to high cross-reactivity of the anti-Tap42 antibody this experiment was inconclusive. Thus our data suggest that the dominant-defective phenotype may be due to competition for substrates or for non-substrate interacting proteins.

#### *Construction of stable cell lines expressing wild type and dominant-defective PP2A -C alleles*

We have also isolated stable fibroblastic cell lines expressing FLAG-tagged mutant and wild-type C $\alpha$  subunits. Interestingly, we consistently see higher expression of the epitope tagged protein in cell lines expressing the H118N mutant construct. Cell lines expressing the FLAG-C $\alpha$ <sup>+</sup> construct show lower transgene expression levels (an estimated three-fold difference on average), and several FLAG-C $\alpha$ <sup>+</sup> cell lines grow slowly. In contrast, the cell lines expressing the H118N mutant at high levels show no obvious growth phenotypes. We are currently quantitating these effects in growth curve assays.

### KEY RESEARCH ACCOMPLISHMENTS

- Construction of dominant-defective alleles of human PP2A-C $\alpha$

- Development of functional epitope-tagged PP2A-C $\alpha$  alleles
- Analysis of the dominant-defective phenotype in yeast cells via co-immunoprecipitation and immunoblotting
- Analysis of A regulatory subunit interaction with mutant and wild-type PP2A-C $\alpha$  alleles in COS cells
- Isolation of stable cell lines expressing epitope-tagged mutant and wild-type PP2A-C $\alpha$  alleles

#### REPORTABLE OUTCOMES

- Lizotte, D., D. D. McManus, H. R. Cohen, and A. DeLong. 1999 Functional expression of human and *Arabidopsis* protein phosphatase 2A in *Saccharomyces cerevisiae* and isolation of dominant-defective mutants. *Gene* 234:35-44
- "Using dominant defective mutants to study protein phosphatase 2A function in vivo" Donna Lizotte, Albert Siryaporn, David McManus, Kimberly Hemond and Alison DeLong; poster presentation at the Brown University MCB Graduate Program Annual Retreat, Sept. 1, 1999.
- "Using dominant defective mutants to study protein phosphatase 2A function in vivo" Donna Lizotte, Albert Siryaporn, David McManus, Kimberly Hemond and Alison DeLong; poster presentation at the Era of Hope BCRP meeting, Atlanta GA, June 8 - 11, 2000.
- Development of stable cell lines expressing native and epitope-tagged wild-type and dominant-defective PP2A-C alleles.

#### CONCLUSIONS

We have shown that *S. cerevisiae* can be used as a rapid in vivo assay system for human PP2A-C activity. Mutation of conserved histidine or arginine residues in the C $\alpha$  sequence confers a strong dominant-defective phenotype. The dominant-defective phenotype does not reflect competition for A or B regulatory subunit binding, and may be due to competition for substrates or other interacting proteins. Our data indicate that even moderate over-expression of wild-type PP2A-C subunit in mammalian cells may lead to reduced growth rates, however, further experiments are required to allow accurate quantitation of this effect.

## REFERENCES

- Campbell, K.S., Auger, K.R., Hemmings, B.A., Roberts, T.M. and Pallas, D.C. (1995) Identification of regions in polyomavirus middle T and small t antigens important for association with protein phosphatase 2A. *J. Virol.* **69**, 3721-3728.
- Cohen, P., Klumpp, S. and Schelling, D.L. (1989) An improved procedure for identifying and quantitating protein phosphatases in mammalian tissues. *FEBS Lett.* **250**, 596-600.
- Di Como, C.J. & Arndt, K.T. (1996) Nutrients, via the Tor proteins, stimulate the association of Tap42 with type 2A phosphatases. *Genes Dev.* **10**, 1904-1916.
- Garbers, C., DeLong, A., Deruère, J., Bernasconi, P. and Söll, D. (1996) A mutation in protein phosphatase 2A regulatory subunit A affects auxin transport in *Arabidopsis*. *EMBO J.* **15**, 2115-2124.
- Pallas, D.C., Shahrik, L.K., Martin, B.L., Jaspers, S., Miller, T.B., Brautigan, D.L. and Roberts, T.M. (1990) Polyoma small and middle T antigens and SV40 small t antigen form stable complexes with protein phosphatase 2A. *Cell* **60**, 167-176.
- Ruediger, R., Roeckel, D., Fait, J., Berqvist, A., Magnusson, F. & Walter, F. (1992) Identification of binding sites on the regulatory A subunit of protein phosphatase 2A for the catalytic C subunit and for the tumor antigens of simian virus 40 and polyomavirus. *Mol. Cell. Biol.* **12**, 4872-4882.
- Sontag, E., Fedorov, S., Kamibayashi, C., Robbins, D., Cobb, M. and Mumby, M. (1993) The interaction of SV40 small tumor antigen with protein phosphatase 2A stimulates the map kinase pathway and induces cell proliferation. *Cell* **75**, 887-897.
- Suganuma, M., Fujiki, H., Suguri, H., Yoshizawa, S., Hirota, M., Nakayasu, M., Ohika, M., Wakamatsu, K., Yamada, K. and Sugimura, T. (1988) Okadaic acid: an additional non-phorbol-12-tetradecanoate-13-acetate-type tumor promoter. *Proc. Natl. Acad. Sci. U.S.A.* **85**, 1768-1771.
- Walter, G. and Mumby, M. (1993) Protein serine/threonine phosphatases and cell transformation. *Biochim. Biophys. Acta* **1155**, 207-226.
- Walter, G., Ruediger, R., Slaughter, C. and Mumby, M. (1990) Association of protein phosphatase 2A with polyoma virus medium tumor antigen. *Proc. Natl. Acad. Sci. USA* **87**, 2521-2525.



Reprinted from

# GENE

---

AN INTERNATIONAL JOURNAL ON  
GENES AND GENOMES

---

Gene 234 (1999) 35–44

[www.elsevier.com/locate/gene](http://www.elsevier.com/locate/gene)

## Functional expression of human and *Arabidopsis* protein phosphatase 2A in *Saccharomyces cerevisiae* and isolation of dominant-defective mutants

D.L. Lizotte<sup>1</sup>, D.D. McManus<sup>1,2</sup>, H.R. Cohen, A. DeLong<sup>\*</sup>

*Department of Molecular Biology, Cell Biology and Biochemistry, Brown University, Providence, RI 02912, USA*



ELSEVIER

# GENE

AN INTERNATIONAL JOURNAL ON  
GENES AND GENOMES

## AIMS AND SCOPE

The journal will publish papers on all structural, functional and evolutionary aspects of genes, chromatin, chromosomes and genomes.

## EDITORIAL POLICY

The overriding criteria for publication are originality, high scientific quality, and interest to a wide audience of readers. Papers outside the scope of the journal, not having the requested scientific criteria of interest or quality, or not having been prepared according to the instructions outlined below, will be sent back to the authors without delay.

The articles published in *Gene* represent the opinions and scientific findings of the authors. Whilst the Publisher and Editors of *Gene* make every effort to ensure the accuracy of all published matter, they can accept no responsibility or liability, collectively or individually, for any erroneous, misleading or unintentionally damaging statements which may appear in the journal. Submission of a paper to *Gene* implies that it is not being published or submitted elsewhere (including symposia volumes or other types of publication).

The Editors assume that all new isolates and mutants described in papers published in *Gene* will be made freely available upon written request to all qualified members of the scientific community. If delays in strain or vector distribution are anticipated or if these are available from sources other than the authors, this should be indicated.

*Gene* requires all new nucleotide and amino acid sequences to be deposited with GenBank before publication [GenBank, Submissions, National Center for Biotechnology Information, Bldg. 38A, Room 8N-803, 8600 Rockville Pike, Bethesda, MD 20894, USA. General Inquiries: e-mail: [info@ncbi.nlm.nih.gov](mailto:info@ncbi.nlm.nih.gov); Tel: +1 (301) 496 2745; Fax: +1 (301) 480 9241. E-mail submission of new sequences: [gb-sub@ncbi.nlm.nih.gov](mailto:gb-sub@ncbi.nlm.nih.gov)]. Obtain and include an accession number with all sequences.

**Publication information.** *Gene* (ISSN 0378-1119). For 1998 volumes 202-220 are scheduled for publication. Subscription prices are available upon request from the Publisher. Subscriptions are accepted on a prepaid basis only and are entered on a calendar year basis. Issues are sent by surface mail except to the following countries where Air delivery via SAL mail is ensured: Argentina, Australia, Brazil, Canada, Hong Kong, India, Israel, Japan, Malaysia, Mexico, New Zealand, Pakistan, PR China, Singapore, South Africa, South Korea, Taiwan, Thailand, USA. For all other countries airmail rates are available upon request. Claims for missing issues should be made within six months of our publication (mailing) date.

**Enquiries concerning manuscripts and proofs:** questions arising after acceptance of the manuscript, especially those relating to proofs, should be directed to:

Issue Manager, *Gene*, Elsevier Science BV, P.O. Box 2759, 1000 CT Amsterdam, The Netherlands; Tel: (+31)-20-4852772; Fax: (+31)-20-4852459; e-mail: [l.koning@elsevier.nl](mailto:l.koning@elsevier.nl)

**Authors in Japan please note:** Upon request, Elsevier Science Japan will provide authors with a list of people who can check and improve the English of their paper (*before submission*). Please contact our Tokyo office: Elsevier Science K.K., 1-9-15 Higashi-Azabu, Minato-ku, Tokyo 106; Tel.: (03) 5561-5032; Fax: (03) 5561-5045.

ALSO AVAILABLE ONLINE:

<http://www.elsevier.nl/locate/gene>

or

<http://www.elsevier.com/locate/gene>

Gene is cited in: Biological Abstracts; Chemical Abstracts; Elsevier BIOBASE/Current Awareness in Biological Sciences; Current Contents (Life Sciences); EMBASE/Excerpta Medica; Index Medicus; Pascal M; Science Citation Index; Reference Update.

**US mailing notice:** *Gene* (ISSN 0378-1119) is published bi-weekly by Elsevier Science B.V. (Molenwerf 1, Postbus 211, 1000 AE Amsterdam, The Netherlands). Annual subscription price in the USA US \$6487.00 (subject to change), including air speed delivery. Periodicals postage paid at Jamaica, NY 11431, USA POSTMASTERS: Send address changes to *Gene*, Publications Expediting, Inc., 200 Meacham Avenue, Elmont, NY 11003. Airfreight and mailing in the USA by Publications Expediting.

Printed in The Netherlands.

**Founded in 1976 by Wacław Szybalski**



# Functional expression of human and *Arabidopsis* protein phosphatase 2A in *Saccharomyces cerevisiae* and isolation of dominant-defective mutants

D.L. Lizotte<sup>1</sup>, D.D. McManus<sup>1,2</sup>, H.R. Cohen, A. DeLong<sup>\*</sup>

Department of Molecular Biology, Cell Biology and Biochemistry, Brown University, Providence, RI 02912, USA

Received 27 December 1998; received in revised form 28 March 1999; accepted 5 May 1999; Received by V. Larionov

## Abstract

Protein phosphatase 2A (PP2A), a heterotrimeric serine/threonine-specific protein phosphatase, comprises a catalytic subunit and two distinct regulatory subunits, A and B. The primary sequence of the catalytic (C) subunit is highly conserved in evolution, and its function has been shown to be essential in yeast, *Drosophila* and mice. In many eukaryotes, the C subunit is encoded by at least two nearly identical genes, impeding conventional loss-of-function genetic analysis. We report here the development of a functional complementation assay in *S. cerevisiae* that has allowed us to isolate dominant-defective alleles of human and *Arabidopsis* C subunit genes. Wild-type human and *Arabidopsis* C subunit genes can complement the lethal phenotype of *S. cerevisiae* PP2A-C mutations. Site-directed mutagenesis was used to create two distinct, catalytically impaired C subunit mutants of the human and *Arabidopsis* genes. In both cases, expression of the mutant subunit in yeast prevented growth, even in the presence of functional C subunit proteins. This dominant growth defect is consistent with a dominant-interfering mode of action. Thus, we have shown that *S. cerevisiae* provides a rapid system for the functional analysis of heterologous PP2A genes, and that two mutations that abrogate phosphatase activity exhibit dominant-defective phenotypes in *S. cerevisiae*. © 1999 Published by Elsevier Science B.V. All rights reserved.

**Keywords:** Catalytic subunit; Dominant-negative; Functional complementation; Mutational analysis

## 1. Introduction

Protein phosphatase 2A (PP2A) is a member of the PPP family of protein serine/threonine phosphatases

Abbreviations: AthC1, AthC3 and AthC5, *Arabidopsis thaliana* PP2A-C genes; CDC55, *S. cerevisiae* PP2A B regulatory subunit gene; Clb2-Cdc28, B-type cyclin/cyclin-dependent kinase; cs, cold-sensitive; HsC $\alpha$  and HsC $\beta$ , human PP2A-C genes; PCR, polymerase chain reaction; PP1, protein phosphatase 1; PP2A, protein phosphatase 2A; PP2A-C, PP2A catalytic subunit; PP2B, protein phosphatase 2B; PPA2, *S. pombe* PP2A-C gene; PPH21 and PPH22, *S. cerevisiae* PP2A-C genes; PPH3, *S. cerevisiae* PP2A-like phosphatase gene; RTS1, *S. cerevisiae* PP2A B' regulatory subunit gene; SDS-PAGE, SDS-polyacrylamide gel electrophoresis; TPD3, *S. cerevisiae* PP2A A regulatory subunit gene; ts, temperature-sensitive; YPD, yeast peptone glucose medium.

<sup>\*</sup> Corresponding author. Tel.: +1-401-863-3888;

fax: +1-401-863-2421.

E-mail address: alison\_delong@brown.edu (A. DeLong)

<sup>1</sup> These authors contributed equally to this work.

<sup>2</sup> Present address: University of Massachusetts Medical Center, Worcester, MA, USA.

(Barford, 1996). Purification of PP2A enzymatic activity from a wide variety of species typically results in the isolation of the PP2A catalytic (C) subunit in a complex with one or more regulatory subunits. A heterotrimeric holoenzyme comprising regulatory A and B subunits bound to the catalytic subunit is thought to predominate in vivo. Interactions between the catalytic and regulatory subunits have been shown to modulate PP2A activity both in vivo and in vitro [reviewed in Mumby and Walter (1993)]. Recent evidence shows that other proteins can bind to the holoenzyme or, in some cases, replace one or both regulatory subunits, resulting in altered PP2A activity [see, for example, Murata et al. (1997)]. The catalytic subunit exhibits a strong sequence similarity to protein phosphatase 1 (PP1) in regions that constitute the putative active site (Goldberg et al., 1995). Although the reaction mechanisms of PP1 and PP2A have not yet been defined, amino acid sequence conservation, crystallographic analysis of the PP1 active site (Goldberg et al., 1995) and biochemical analysis of the bacteriophage  $\lambda$  phosphatase (Zhuo et al., 1994) have

|        |             |            |            |            |            |            |     |
|--------|-------------|------------|------------|------------|------------|------------|-----|
|        | 1           |            |            |            |            |            | 60  |
| Pph21p | mdtdldvpmq  | davteqltpt | vsemdlnnn  | ssdnna.... | ....eefsvd | dlkpgssgia |     |
| Pph22p | mdmeiddpmh  | gsdedqlspt | ldedmsddg  | knntkarsnd | edtdeeledf | nfkpgssgia |     |
|        | 61          |            |            |            |            |            | 120 |
| Pph21p | dhksskpllel | nntningLDq | wIEhLskCep | LsEddVarLC | kmAvdVLqfE | eNVkpInvPV |     |
| Pph22p | dhksskplkl  | tntningLDq | wIEhLskCep | LsEddVarLC | kmAvdVLqfE | eNVkpInvPV |     |
| AthC1  | .....       | .mplngdLDr | qIEqLmeCkp | LgEadVkiLC | dqAkaILVeE | yNVqpVkcPV |     |
| HsCα   | .....md     | ekvftkeLDq | wIEqLneCkq | LsEsqVksLC | ekAkeILtke | sNVqeVrcPV |     |
| HsCβ   | .....md     | dkaftkeLDq | wVEqLneCkq | LnEnqVrtLC | ekAkeILtke | sNVqeVrcPV |     |
| CONSEN | -----       | -----LD-   | -IE-L--C-- | L-E--V--LC | --A--IL--E | -NV--V--PV |     |
|        | 1           |            |            |            | K†         | 52         |     |
|        | 121         |            |            |            |            |            | 180 |
| Pph21p | TICGDVHGQF  | hDLlELFkIG | GpcPDTNYLF | MGDYVDRGY  | SVETVsylVA | MKVRYPHrIT |     |
| Pph22p | TICGDVHGQF  | hDLlELFkIG | GpcPDTNYLF | MGDYVDRGY  | SVETVsylVA | MKVRYPHrIT |     |
| AthC1  | TVCGDIHGQF  | yDLlELFrIG | GnapDTNYLF | MGDYVDRGY  | SVETVsllVA | LKVRYPHrIT |     |
| HsCα   | TVCGDVHGQF  | hDLmELFrIG | GksPDTNYLF | MGDYVDRGY  | SVETVtlLVA | LKVRYPHrIT |     |
| HsCβ   | TVCGDVHGQF  | hDLmELFrIG | GksPDTNYLF | MGDYVDRGY  | SVETVtlLVA | LKVRYPHrIT |     |
| CONSEN | TVCGDVHGQF  | -DL-ELF-IG | G--PDTNYLF | MGDYVDRGY  | SVETV--LVA | LKVRYPHrIT |     |
|        | 53          |            |            | A          |            | 112        |     |
|        | 181         |            |            |            |            |            | 240 |
| Pph21p | ILRGNHESRQ  | ITQVYGFYDE | CLRKYGSANV | WKmFTDLFDY | FPtALVDnk  | IFCLHGGLSP |     |
| Pph22p | ILRGNHESRQ  | ITQVYGFYDE | CLRKYGSANV | WKmFTDLFDY | FPvTALVDnk | IFCLHGGLSP |     |
| AthC1  | ILRGNHESRQ  | ITQVYGFYDE | CLRKYGSANV | WKyFTDLFDY | LPtALIESq  | VFCLHGGLSP |     |
| HsCα   | ILRGNHESRQ  | ITQVYGFYDE | CLRKYGSANV | WKyFTDLFDY | LPtALVDgq  | IFCLHGGLSP |     |
| HsCβ   | ILRGNHESRQ  | ITQVYGFYDE | CLRKYGSANV | WKyFTDLFDY | LPtALVDgq  | IFCLHGGLSP |     |
| CONSEN | ILRGNHESRQ  | ITQVYGFYDE | CLRKYG-ANV | WK-FTDLFDY | LP-TALVD-- | IFCLHGGLSP |     |
|        | 113         | N          |            |            | S†         | H†         |     |
|        | 241         |            |            |            |            |            | 300 |
| Pph21p | miETIdqvRe  | LnRiQEVPhE | GPMCDLLWSD | PDDRgGWGIS | PRGAGFTFGQ | DVseqFNHtN |     |
| Pph22p | miETIdqvVRd | LnRiQEVPhE | GPMCDLLWSD | PDDRgGWGIS | PRGAGFTFGQ | DiseqFNHtN |     |
| AthC1  | sLDtLDnIRs  | LdRiQEVPhE | GPMCDLLWSD | PDDRcGWGIS | PRGAGYTFGQ | DIatqFNHnN |     |
| HsCα   | siDTLDhIRa  | LdRiQEVPhE | GPMCDLLWSD | PDDRgGWGIS | PRGAGYTFGQ | DIsetFNHaN |     |
| HsCβ   | siDTLDhIRa  | LdRiQEVPhE | GPMCDLLWSD | PDDRgGWGIS | PRGAGYTFGQ | DIsetFNHaN |     |
| CONSEN | --DT-D-IR-  | L-R-QEVPhE | GPMCDLLWSD | PDDR-GWGIS | PRGAGYTFGQ | DI---FNH-N |     |
|        | 173         |            |            |            |            | 232        |     |
|        | 301         |            |            |            |            |            | 360 |
| Pph21p | dLsLIaRAHQ  | LVMEGYawsh | qqNVVTIFSA | PNYCYRCGNq | AAIMEvdEnh | nrqFLQYDPs |     |
| Pph22p | dLsLIaRAHQ  | LVMEGYswsh | qqNVVTIFSA | PNYCYRCGNq | AAIMEvdEnh | nrqFLQYDPs |     |
| AthC1  | gLsLIaRAHQ  | LVMEGYnwsh | ekNVVTIFSA | PNYCYRCGNm | AAILEigEkm | eqnFLQFDPa |     |
| HsCα   | gLtLVsRAHQ  | LVMEGYnwsh | drNVVTIFSA | PNYCYRCGNq | AAIMEldDt1 | kysFLQFDPa |     |
| HsCβ   | gLtLVsRAHQ  | LVMEGYnwsh | drNVVTIFSA | PNYCYRCGNq | AAIMEldDt1 | kysFLQFDPa |     |
| CONSEN | -L-LI-RAHQ  | LVMEGY-W-- | --NVVTIFSA | PNYCYRCGN- | AAIME--E-- | ---FLQFDP- |     |
|        | 233         | Q*         |            | ****       |            | 292        |     |
|        | 361         | 377        |            |            |            |            |     |
| Pph21p | vRpgEPsvsR  | kTPDYFL    |            |            |            |            |     |
| Pph22p | vRpgEPtvTR  | kTPDYFL    |            |            |            |            |     |
| AthC1  | pRqvEPdtTR  | kTPDYFL    |            |            |            |            |     |
| HsCα   | pRrgEPhvTR  | rTPDYFL    |            |            |            |            |     |
| HsCβ   | pRrgEPhvTR  | rTPDYFL    |            |            |            |            |     |
| CONSEN | -R--EP--TR  | -TPDYFL    |            |            |            |            |     |
|        | 293         | KVTR       | RTPDYFL    |            |            |            |     |

identified residues required for metal ion coordination, substrate binding, and catalysis (see Fig. 1).

The amino acid sequences of C and A subunits are highly conserved in eukaryotes as distantly related as *Arabidopsis thaliana*, humans, and yeast. Conservation of the A subunit function has been demonstrated previously by complementing a *S. cerevisiae* A subunit mutant with an *Arabidopsis* A subunit gene (Garbers et al., 1996). In the animals studied to date, both the C and A subunits are expressed ubiquitously and appear to be encoded by small and fairly homogeneous gene families. In plants, however, A and C subunit gene families are larger; the *Arabidopsis* genome encodes at least five C subunits and three A subunits [reviewed in Smith and Walker (1996) and Stamey and Rundle (1996)]. In both plants and animals, different B subunit isoforms are encoded by two or more unrelated gene families, some of which are expressed in a tissue-specific manner. Thus, the more variable B subunit may dictate substrate specificity (Mayer-Jaekel and Hemmings, 1994), a hypothesis supported by data showing that the tumor antigens of DNA tumor viruses replace B subunits in PP2A complexes and alter the enzyme's catalytic properties (Yang et al., 1991; Cayla et al., 1993; Sontag et al., 1993).

PP2A activity has been shown to play a negative role in growth control in systems such as *Xenopus*, *S. pombe*, and *Drosophila* (Lee et al., 1991; Kinoshita et al., 1993; Mayer-Jaekel et al., 1994). In *S. cerevisiae*, however, PP2A is required for entry into mitosis (Lin and Arndt, 1995). Two PP2A-C genes, designated *PPH21* and *PPH22*, have been identified in *S. cerevisiae*. Inactivation of both genes is detrimental but not lethal unless a gene encoding a PP2A-like phosphatase, *PPH3*, is also mutated (Sneddon et al., 1990; Ronne et al., 1991; Lin and Arndt, 1995). Cells carrying a temperature-sensitive (ts) *pph21-102* allele in a *pph21 pph22 pph3* null background retain viability but exhibit G2 cell-cycle arrest, reduced B-type cyclin/cyclin-dependent (Clb2-Cdc28) kinase activity and abnormal bud morphologies at the non-permissive temperature (Lin and Arndt, 1995). The PP2A-A subunit is encoded by a single gene, *TPD3*, and *tpd3* null mutants exhibit a Ts<sup>-</sup> phenotype (van Zyl et al., 1992). Two B subunits have been identified;

mutants in the 55-kDa B subunit gene *CDC55* are cold-sensitive (cs), whereas mutants in the 56-kDa B' subunit gene *RTS1* are ts, cs and ethanol-hypersensitive (Healy et al., 1991; Shu et al., 1997). It is unclear why the C subunit is essential, whereas the A and B regulatory subunits are only conditionally required.

The fact that each PP2A subunit is encoded by a gene family in many eukaryotes presents a significant impediment to conventional genetic analysis, and suggested to us that the isolation of dominant PP2A mutants might provide an alternative to the isolation of mutations in individual gene family members followed by the generation of multiply mutant stocks. Taking advantage of the high sequence conservation exhibited by PP2A genes and the availability of PP2A mutant strains of yeast, we have developed a functional assay for the isolation and characterization of dominant-defective alleles of PP2A catalytic subunit genes. We have demonstrated the utility of this system by the construction of dominant-defective PP2A-C mutants from *Arabidopsis thaliana* and from humans.

## 2. Materials and methods

### 2.1. Yeast strains and plasmid constructs

The yeast strains used in this work are listed in Table 1. The yeast growth media and genetic techniques were as described by Ausubel et al. (1992). After lithium acetate transformation, yeast transformants were selected at 25°C on synthetic complete glucose (SCD) medium lacking tryptophan and uracil. Segregants that had lost either the *pph21-102*/YCp50 construct or the heterologous C subunit construct were isolated by growing cells non-selectively (on YPD or YPGalactose medium at 25°C), followed by replica-patching single colonies onto selective (SCD or SCGalactose) medium with or without uracil or tryptophan. Segregants were colony-purified and their phenotypes verified by restreaking on the appropriate media.

All *GAL* constructs used in this work are derivatives of either YCp22GAL or YEpl12GAL (shown in Table 1 as *pGAL/TRP/CEN* and *pGAL/TRP/2μ*, respectively),

Fig. 1. Comparison of PP2A catalytic subunit amino acid sequences. Residues likely to be critical active site residues are underlined (Goldberg et al., 1995). The substitutions found in catalytic subunit mutants are shown below the consensus sequence, with bold letters indicating mutations generated in this work. Also shown are *pph21* and *pph22* mutations conferring temperature sensitivity in *S. cerevisiae* (†: Lin and Arndt, 1995; Evans and Stark, 1997) and a *ppa2* mutation conferring cold sensitivity in *S. pombe* (\*: Kinoshita et al., 1990). The sites of several mutations conferring okadaic acid resistance are also indicated (^: Shima et al., 1994; Kaneko et al., 1995). Numbering above each sequence block refers to the amino acid sequence of Pph22p, and italicized numbering below the sequence blocks refers to the sequence of the human gene products. Two additional *Arabidopsis* catalytic subunit sequences (AthC3 and AthC5) were included in the sequence comparison, but are not shown in the figure. Residues conserved in six of the seven sequences compared are shown in the consensus sequence (CONSEN). Sequence identities and conservative substitutions are represented by upper-case letters, and non-conservative substitutions are shown in lower-case letters. The sequence of the peptide used to raise the anti-PP2A-C antibody used in these studies is shown in italics under the carboxy-terminal end of the consensus sequence. GenBank Accession Nos: PPh21p, X56261; Pph22p, X56262; AthC1, M69732; AthC3, M96841; AthC5, U39568; HsCα, M36951; HsCβ, X12656. The alignment and consensus sequence were generated using the PILEUP and PRETTY algorithms of the University of Wisconsin GCG package.

Table 1  
Strain list

|        |  |                      |
|--------|--|----------------------|
| W303   | <i>MATa ura3-1 leu2-3,112 his3-11,15 trp1-1 ade2-1 can1-100</i>                                    | K. Arndt             |
| CY3007 | <i>MATa pph22::HIS3 pph21::HIS3 pph3::LEU2 [pph21-102/URA/CEN] W303</i>                            | Lin and Arndt (1995) |
| DLY6   | <i>MATa pph22::HIS3 pph21::HIS3 pph3::LEU2 LEU2::ADH:AthC1 kan<sup>r</sup> [pph21-102/URA/CEN]</i> | This work            |
| DLY8   | <i>CY3007 [ADH:AthC1/TRP/CEN]</i>  | This work            |
| DLY9   | <i>CY3007 [ADH:AthC1-antisense/TRP/CEN]</i>  | This work            |
| DLY10  | <i>CY3007 [pGAL/TRP/CEN]</i>   | This work            |
| DLY11  | <i>CY3007 [pGAL/TRP/2μ]</i>  | This work            |
| DLY12  | <i>CY3007 [GAL10:AthC1/TRP/2μ]</i>   | This work            |
| DLY13  | <i>CY3007 [GAL10:AthC1-H115N/TRP/2μ]</i>   | This work            |
| DLY14  | <i>CY3007 [GAL10:AthC1-R86A/TRP/2μ]</i>  | This work            |
| DLY16  | <i>CY3007 [GAL1:HsCα/TRP/2μ]</i>   | This work            |
| DLY17  | <i>CY3007 [GAL1:HsCα/TRP/CEN]</i>  | This work            |
| DLY18  | <i>CY3007 [GAL1:HsCβ/TRP/CEN]</i>  | This work            |
| DLY19  | <i>CY3007 [GAL1:HsCα-H118N/TRP/CEN]</i>  | This work            |
| DLY20  | <i>CY3007 [GAL1:HsCα-R89A/TRP/CEN]</i>   | This work            |
| DLY21  | <i>DLY6 [pGAL/TRP/2μ]</i>  | This work            |
| DLY22  | <i>DLY6 [GAL10:AthC1/TRP/2μ]</i>   | This work            |
| DLY23  | <i>DLY6 [GAL10:AthC1-H115N/TRP/2μ]</i>   | This work            |
| DLY24  | <i>DLY6 [pGAL/TRP/CEN]</i>   | This work            |
| DLY25  | <i>DLY6 [GAL10:AthC1/TRP/CEN]</i>  | This work            |
| DLY26  | <i>DLY6 [GAL10:AthC1-H115N/TRP/CEN]</i>  | This work            |

isogenic yeast expression vectors carrying the *GAL1/GAL10* bidirectional promoter, a *TRP1* selectable marker, and a *CEN* or 2μ circle origin of replication, respectively (Pitluk et al., 1995). Plasmid pADL302 (*GAL10:AthC1/TRP/2μ*) was constructed by subcloning the *AthC1* coding sequence from pPP2A-1 (a kind gift of Sabine Rundle) into the *EcoRI* site of YEpl12GAL. Plasmids pDMC2 and pDMC16 (*GAL1:HsCα/TRP/CEN* and *GAL1:HsCα/TRP/2μ*) were constructed as follows: the *HsCα* coding sequence was amplified from plasmid pUC.HPP2ACαRM1 (a kind gift of N. Andjelkovic and B. Hemmings) and given *Bam*HI cloning ends via polymerase chain reaction (PCR), followed by cleavage with *Bam*HI and ligation into *Bam*HI-cleaved YCp22GAL and YEpl12GAL, respectively. The primers used were 5' GCGGATCCTTACAGGAAGTAGTCTGGGG 3' and 5' CGGGATCCATGGACGAGAAGGTGTTACCAAGG 3'. Similarly, plasmid pDMC4 (*GAL1:HsCβ/TRP/CEN*) was constructed by PCR amplification of the *HsCβ* coding sequence from plasmid pTZ18U.HFP6A2Cβ (a kind gift of N. Andjelkovic and B. Hemmings), followed by *Bam*III subcloning into YCp22GAL, using PCR primers 5' GCGGATCCTTATAGGAAGTAGTCTGGGG 3' and 5' CGGGATCCATGGACGACAAGGCGTTCACCAAGG 3'.

To make the *ADH:AthC1* constructs, a 2-kb *Bam*HI fragment carrying the *ADH* promoter and terminator was subcloned from pAAH5 (Ammerer, 1983) into pTZ19 (US Biochemical). The *AthC1* coding sequence was ligated into the unique *Hind*III site between the *ADH* promoter and terminator. To make plasmids pADL316 (*ADH:AthC1/TRP/CEN*) and pADL317 (*ADH:AthC1 antisense/TRP/CEN*), sense and antisense

*ADH:AthC1* fusions were subcloned as *Sph*I fragments into YCplac22 (Gietz and Sugino, 1988). To chromosomally integrate the *ADH:AthC1* construct, the pADL316 *Sph*I fragment was cloned into the integrating plasmid YIplac128 (Gietz and Sugino, 1988), followed by addition of a G418 resistance cassette (*Sma*I–*Sac*I fragment) from pFA6-KanMX4 (Wach et al., 1994). The resulting plasmid (pADL340) was linearized with *Afl*II and transformed into strain CY3007, with selection for G418 resistance.

## 2.2. Site-directed mutagenesis

Mutants were generated using an oligonucleotide-mediated mutagenesis protocol (Deng and Nickoloff, 1992). The following mutagenic oligonucleotide primers were used. oAthC1-H115N: 5' CTGACGGCTCTCA-TTATTCCTCGCAGG 3'; oAthC1-R86A: 5' CAG-AATAGTAGCCAGCATCTACGTAATCTCCCATG-AAG 3'; oHsCα-H118N: 5' TTCTTCGAGGGAATA-ATGAGAGCAGACAG 3'; oHsCα-R89A: 5' GGG-AGATTATGTTGATGCAGGATATTATTTCAGTT 3'.

## 2.3. Immunoblot analysis

Standard SDS–PAGE and immunoblotting protocols were followed (Ausubel et al., 1992). The polyclonal anti-PP2A-C antibody used was raised against a conserved carboxy-terminal peptide (KVRRTPDYFL), and was a kind gift of T. Stukenberg and M. Kirschner. Peroxidase-conjugated goat anti-rabbit secondary antibody (Jackson ImmunoResearch Laboratories) was detected using a standard chemiluminescence protocol.

### 3. Results

#### 3.1. *Arabidopsis* and human PP2A-C clones complement a yeast PP2A mutant

We established a functional complementation assay for PP2A activity in *S. cerevisiae*. Strain CY3007 carries null alleles in both PP2A-C genes and in the *PPH3* gene, which encodes a related phosphatase; lethality of this triple mutation is rescued by a low-copy-number plasmid bearing a *ts pph21* allele (*pph21-102*; Lin and Arndt, 1995). CY3007 cells grow at 25 or 30°C, but not at 35°C. We cloned *Arabidopsis* and human cDNAs encoding PP2A-C subunits into yeast expression vectors and tested their ability to complement the temperature sensitivity of strain CY3007. When expressed from the constitutive alcohol dehydrogenase (ADH) promoter and carried on a low-copy-number plasmid, the *Arabidopsis* AthC1 cDNA (see Fig. 1; Ariño et al., 1993) complemented the *Ts*<sup>−</sup> phenotype of CY3007, indicating that the *Arabidopsis* PP2A-C subunit is functional in yeast (Fig. 2). When cloned in the antisense orientation, the AthC1 cDNA failed to complement in CY3007 (Fig. 2).

To exclude the possibility that the yeast *ts* allele contributed to complementation, we grew cells carrying the ADH-AthC1 plasmid non-selectively and screened for tryptophan or uracil auxotrophy to identify segregants that had lost either plasmid. Loss of the ADH-AthC1 plasmid conferred a *Trp*<sup>−</sup> phenotype and restored the parental *Ts*<sup>−</sup> phenotype. Loss of the *pph21-102* plasmid conferred a *Ura*<sup>−</sup> phenotype, but the resulting cells still grew at 35°C, as well as 25 and 30°C (Fig. 2). Thus, the ADH-AthC1 plasmid is necessary and sufficient for complementing the *Ts*<sup>−</sup> phenotype,

and is capable of supporting normal growth in the absence of any yeast PP2A-C protein.

Expression of either the AthC1 or the human HsCα cDNA (see Fig. 1; Stone et al., 1988) under the control of a galactose-inducible *GAL* promoter also complemented the *Ts*<sup>−</sup> phenotype of CY3007. Cells expressing either the *Arabidopsis* or human cDNA grew at 25°C on both glucose- and galactose-containing media and at 35°C on galactose, but did not grow on glucose at 35°C (Fig. 3A and B). The sufficiency of the *GAL*:AthC1 construct was tested by segregating either the *GAL*:AthC1 or YCp50 *pph21-102* plasmid. Again, loss of the plasmid carrying the *Arabidopsis* C1 gene restored the parental *Ts*<sup>−</sup> phenotype. Loss of the *pph21-102* plasmid conferred a galactose-dependent phenotype; segregants were viable at 25, 30 and 35°C only on galactose-containing media (data not shown). Surprisingly, the human HsCβ gene (see Fig. 1; Hemmings et al., 1988) did not provide complementing PP2A activity in yeast (Fig. 3B). Failure to complement was not due to poor expression of the HsCβ protein since similar amounts of the HsCα and β subunit proteins were detected in galactose-grown cells (Fig. 4A). Thus, expression of either the AthC1 or HsCα (but not HsCβ) catalytic subunit gene provides the PP2A function in yeast cells.

We predict that cells expressing a catalytically active C subunit that was unable to interact with the endogenous yeast regulatory subunits (e.g. *TPD3* or *RTS1*) would exhibit phenotypes characteristic of loss-of-function mutations in those genes. Both *tpd3* and *rts1* mutations prevent growth at 37°C (van Zyl et al., 1992; Shu et al., 1997); therefore, we tested the ability of cells expressing human and *Arabidopsis* PP2A-C subunits to grow at 37°C. Cells carrying a *GAL*:HsCα or *GAL*:AthC1 construct were viable at 37°C, suggesting that these heterologous C subunits functionally interact with the yeast A and B' regulatory subunits (Fig. 3C). Cells carrying the *GAL*:AthC1 construct formed small colonies at this temperature. This small colony phenotype may indicate that the AthC1 protein has lower affinity for Tpd3p than does HsCα, however, it is also possible that the catalytic activity of AthC1 is reduced at 37°C, a temperature above the normal growth range of *Arabidopsis*.

High-level over-expression of PP2A-C mRNA has been shown to be toxic in *S. cerevisiae*, but cells expressing the *PPH22* gene under control of the *GAL1* promoter grow at a normal rate (Ronne et al., 1991). We have not observed any strong negative growth effects of *GAL*-driven expression of wild-type human or *Arabidopsis* PP2A-C genes. In fact, under permissive conditions, cells carrying a high-copy-number *GAL*:C construct grow slightly faster in liquid media (c. 2.5 h doubling time) than do cells carrying the vector alone (c. 3 h doubling time).




Fig. 2. Growth of strains carrying ADH-AthC1 plasmids. Yeast cells from single colonies were patched on duplicate plates containing YPD medium and grown at the temperatures shown. The PP2A-C alleles carried by cells in each set of patches are shown (top): *Ts*<sup>+</sup>, parental strain W303; *ts*, temperature-sensitive mutant strain CY3007; *ts* + C1, DLY8; *ts* + anti-C1, DLY9; (*ts* + C1) − *ts*, *Ura*<sup>−</sup> derivatives of DLY8; (*ts* + C1) − C1, *Trp*<sup>−</sup> derivatives of DLY8. Similar results were obtained in streak-out tests of growth of these strains.




Fig. 3. Growth of yeast expressing heterologous C subunits under *GAL* promoter control. CY3007 transformants carrying a *GAL10:AthC1* construct (AthC1; DLY12) or an empty vector (–; DLY11) were streaked on duplicate plates containing selective glucose and galactose medium and incubated at the temperature indicated (A). CY3007 transformants carrying an empty vector (sector 1; DLY11) or low copy (sector 2; DLY17) or high copy (sector 4; DLY16) *GAL1:HsCα*, or low copy *GAL1:HsCβ* (sector 3; DLY18) were streaked on duplicate plates containing selective glucose and galactose medium and incubated at the temperature indicated (B). CY3007 transformants carrying an empty vector (quadrant 5; DLY11), low copy (quadrant 6; DLY17) or high copy (quadrant 7; DLY16) *GAL1:HsCα*, or high-copy *GAL10:AthC1* (quadrant 8; DLY12) were streaked on selective galactose medium and incubated at the temperature indicated (C).

### 3.2. Isolation of dominant defective alleles of *PP2A-C*

We hypothesized that mutation of an amino acid residue involved in catalysis would produce a dominant-




Fig. 4. *GAL*-driven expression of wild-type and dominant mutant C subunits in yeast. Cells carrying an empty vector (–; DLY10), and low-copy *GAL1:HsCα* (DLY17) and *GAL1:HsCβ* (DLY18) constructs were grown in galactose, harvested, lysed, and total soluble proteins were extracted and subjected to immunoblot analysis (A). Wild-type and mutant AthC1 proteins were detected after induction of the *GAL10* promoter (B). Cells carrying an empty vector (–; DLY11), *GAL10:AthC1* (DLY12) or *GAL10:AthC1-H115N* (DLY13) were grown in selective raffinose medium, diluted into selective galactose medium at time 0, and samples were harvested after 1 and 3 h, as indicated. Cells were lysed, and proteins were extracted and analyzed as described above. The positions of the yeast C subunit (Pph21p) and the human (HsC) and *Arabidopsis* (AthC) C subunit proteins are indicated on the right of each panel.

defective PP2A-C allele if substrate binding and/or subunit interactions were unaffected by the lesion. We chose a putative active site histidine (H115 in the *Arabidopsis* gene and H118 in the human sequence; see Fig. 1) and mutated this residue to asparagine using site-directed mutagenesis (see Section 2.2). Mutations at the analogous position abrogate catalytic activity but not substrate binding in calcineurin (PP2B) and  $\lambda$  phosphatase (Zhuo et al., 1994; Mertz et al., 1997), two enzymes that exhibit a close similarity to PP2A in putative active site sequences. To test the activity of the mutant proteins, we expressed the mutant alleles under control of *GAL* promoters in strain CY3007 and assayed their effects on growth (Fig. 5). Cells carrying the *GAL:AthC1-H115N* construct were unable to grow on either galactose or glucose at 35°C (Fig. 5A), indicating that the mutant enzyme is defective and provides no catalytic activity at 35°C. More importantly, cells carrying this construct were unable to grow on galactose at 25°C, the permissive temperature for the *pph21-102* allele (Fig. 5A) or at 30°C (data not shown). Thus, the defective phenotype of the H115N allele is dominant over the ts yeast PP2A allele and prevents growth at all temperatures tested. We have obtained the same results with the H118N mutant of the human C $\alpha$  subunit (Fig. 6B and data not shown).

To test the dominance of the H115N allele with






Fig. 5. Growth of yeast cells carrying AthC1-H115N constructs, CY3007 (A) and DLY6 (B and C) transformants carrying an empty vector (–), a *GAL10*:AthC1 construct (+) or a *GAL10*:AthC1-H115N construct (H115N) were streaked on duplicate plates containing selective glucose and galactose medium and incubated at the temperature indicated. Plasmid carried high-copy number (2 $\mu$  circle; A and B) or low-copy-number (centromeric; C) origins of replication. Strains used: DLY11, DLY12, DLY13 (A); DLY21, DLY22, DLY23 (B); DLY24, DLY25, DLY26 (C). Similar results were obtained in streak-out tests of growth of these strains.

respect to the homologous wild-type gene, we chromosomally integrated a wild-type AthC1 gene under control of the ADH promoter. The resulting strain, DLY6, grew well at 30 or 35°C on glucose- or galactose-containing media (Fig. 5B), indicating that the integrated ADH-AthC1 fusion supplies PP2A activity. DLY6 cells carrying the *GAL*:AthC1-H115N construct on a high-copy-number plasmid grew on glucose-containing media at either 30 or 35°C, but failed to grow on galactose at either temperature (Fig. 5B). Thus, the H115N allele is dominant over a wild-type AthC1 allele. The expression




Fig. 6. Growth of yeast cells carrying dominant PP2A-C mutants, CY3007 transformants carrying wild-type (+), and mutant (H115N and R86A) *GAL10*:AthC1 constructs (A: strains DLY12, DLY13 and DLY14) or wild-type and mutant (H118N) *GAL1*:HsC $\alpha$  constructs or wild-type *GAL1*:HsC $\beta$  (B: strains DLY17, DLY19 and DLY18) were streaked on duplicate plates containing selective galactose medium and incubated at the temperature indicated. Similar results were obtained in streak-out tests of growth of these strains.

level of the mutant protein influenced the strength of the dominant-defective phenotype, as a low-copy-number AthC1-H115N plasmid caused a weaker, but still clearly detectable, inhibition of growth (Fig. 5C). This copy number effect is consistent with the dominant-interfering activity of the mutant proteins.

To show that both the mutant and endogenous PP2A-C subunits are expressed in galactose-grown cells, we used immunoblotting to assay PP2A-C subunit levels after galactose induction of cells carrying mutant and wild-type *GAL10*:AthC1 constructs (Fig. 4B). We have obtained the same results with wild-type and H118N mutant human HsC $\alpha$  (data not shown). Wild-type and mutant PP2A-C proteins accumulate to comparable levels after galactose induction, and appear equally stable over a 9-h time course (data not shown).

To determine whether other active site mutations also would confer a dominant-defective phenotype, we introduced a lesion at a position thought to be essential for binding of the phosphorylated substrate. Conversion of an active site arginine to alanine in the  $\lambda$  phosphatase causes a 20-fold decrease in binding of a synthetic substrate and a 500-fold catalytic defect (Zhuo et al., 1994). We introduced the same mutation (R86A in the *Arabidopsis* protein and R89A in the human protein; see Fig. 1) into our wild-type *GAL*:PP2A-C constructs

and tested the activity of the mutant constructs as described above. Cells carrying AthC1-R86A failed to grow on either glucose or galactose at 35°C (Fig. 6 and data not shown), showing that, like the H115N allele, this mutant provides no complementing phosphatase activity at 35°C. Expression of the R86A allele also blocked growth of CY3007 cells at 25°C (Fig. 6A). The R86A mutant protein accumulates to levels comparable to those of the wild-type and H118N proteins after galactose induction (data not shown). These data show that the R86A mutation is also dominant-defective, and we have obtained similar results with the human HsC $\alpha$ -R89A mutant (data not shown). In contrast, the HsC $\beta$  provides no complementing phosphatase activity at 35°C and does not exhibit a dominant-defective phenotype (Fig. 6B).

#### 4. Discussion

We have used yeast to screen for dominant-defective mutants of the PP2A catalytic subunit. We have shown that both *Arabidopsis* and human PP2A genes complement a ts mutation in a yeast PP2A-C gene. Constitutive or inducible expression of a heterologous PP2A-C subunit gene is sufficient to allow growth of yeast cells in the absence of any yeast PP2A-C protein. Site-directed mutagenesis of the *Arabidopsis* and human PP2A-C genes yielded catalytically inactive alleles that exhibited dominant phenotypes. Expression of the dominant-defective phenotype was influenced by copy number, consistent with a dominant-interfering effect of the mutant gene products.

The robust growth of yeast cells in which PP2A function is supplied by a heterologous C subunit protein indicates that sequences required for specific and functional interaction with essential substrates are conserved between the yeast, *Arabidopsis* and human PP2A-C subunits. The predicted amino acid sequences of the C subunit gene products show a very high degree of similarity, 80–90% over the length of the *Arabidopsis* and human sequences. Both yeast proteins carry amino-terminal extensions that are approximately 70 amino acids in length and serve unknown functions. Our work shows that the extensions are not required for essential PP2A functions in *S. cerevisiae* since the *Arabidopsis* and human clones are expressed from their native start codons. Complementation also reveals that critical recognition features of the substrate proteins are conserved. The full range of PP2A substrates required for growth of yeast cells is not known, but the available evidence suggests that essential substrates include proteins involved in bud morphogenesis, cytoskeletal organization and entry into mitosis (Ronne et al., 1991; Lin and Arndt, 1995; Evans and Stark, 1997). Finally, we tested the competence of the human and *Arabidopsis* C

subunits to interact with the endogenous yeast regulatory subunits by assaying for growth at the restrictive temperature for the *tpd3* (A subunit) and *rts1* (B' subunit) mutants. Our data support the hypothesis that sequences required for interaction with regulatory subunits also are conserved in these heterologous proteins, consistent with earlier observations that an *Arabidopsis* A regulatory subunit complements a *tpd3* null allele in cells expressing wild-type yeast C subunits (Garbers et al., 1996).

Although the human HsC $\alpha$  subunit supports growth of yeast, the HsC $\beta$  subunit does not provide complementing PP2A activity in *S. cerevisiae*. The non-complementing phenotype of the C $\beta$  subunit construct is surprising because only eight residues differ in the predicted amino acid sequences of the C $\alpha$  and C $\beta$  subunits. Most of the substitutions are conservative changes in sequences not known to be essential for PP2A function. However, it has recently been shown that mice homozygous for a PP2A-C $\alpha$  null mutation die during embryogenesis, despite significant levels of C $\beta$  subunit expression (Götz et al., 1998). Thus, failure of the C $\beta$  subunit to complement in yeast may reflect functional specificity in vivo.

We used site-directed mutagenesis to create two dominant-defective alleles of PP2A (H115N or H118N and R86A or R89A). Target residues for mutagenesis were chosen on the basis of putative active site function (Goldberg et al., 1995) and biochemical analysis of mutants of the bacteriophage  $\lambda$  phosphatase (Zhuo et al., 1994). Both mutants were inactive in our complementation assay, indicating loss of catalytic activity. The H115N/H118N mutant exhibited a strong dominant-defective phenotype, whereas the R86A/R89A mutant displayed a slightly weaker dominant-defective phenotype. Preliminary microscopic characterization of cells expressing the H115N/H118N allele reveals abnormal bud morphologies similar to those exhibited by PP2A-depleted cells (Ronne et al., 1991) or *cde55* mutant cells (Healy et al., 1991), consistent with the hypothesis that these cells undergo growth arrest due to loss of PP2A function (D. Lizotte, A. DeLong, unpublished). The H115N mutant of AthC1 was dominant in the presence of a constitutively expressed wild-type copy of AthC1, showing that the dominant-defective phenotype was not specific to the Pph21-102p protein. A partially dominant mutant of *S. pombe* PP2A-C has been isolated previously. A high dose of this cold-sensitive (cs) *ppa2* allele was shown to arrest growth under non-permissive conditions, and to retard growth and reduce cell size under permissive conditions (Kinoshita et al., 1990). The basis of the mutant's partial dominance is unknown. However, expression of the cs allele increased okadaic acid sensitivity mildly, suggesting a dominant-interfering mechanism (Kinoshita et al., 1993).

The severity of the dominant growth defect phenotype was clearly affected by the copy number of the mutant and wild-type genes, consistent with the hypothesis that the dominant-defective enzyme competes with the functional C subunits for substrates or regulatory subunits. Either a substrate sequestration or a regulatory subunit titration model could account for the data we have presented here. The HsC $\alpha$ -H118N mutant has been independently constructed and shown to be catalytically inactive despite normal substrate binding characteristics (H. Chung, D. Brautigan, pers. commun.). Equivalent mutants of  $\lambda$  phosphatase and calcineurin display the same biochemical characteristics, but have not been assayed for phenotypes in vivo (Zhuo et al., 1994; Mertz et al., 1997). In contrast, a  $\lambda$  phosphatase mutant equivalent to the R86A/R89A mutant constructed here exhibits a 20-fold decrease in substrate binding and a catalytic defect less severe than that of the histidine mutant (Zhuo et al., 1994). The weaker phenotype of the R86A/R89A mutant may indicate that substrate binding contributes to, but is not essential for, the dominant phenotype. A biochemical analysis will be required to resolve these questions. Interestingly, the HsC $\beta$  subunit does not provide complementing PP2A activity, but is not dominant-defective. This result suggests that the factors required for functional complementation also are required for the dominant-defective phenotype, and argues against the hypothesis that any non-functional C subunit may exert a dominant effect. The partially dominant mutation conferring cold sensitivity on *S. pombe* PP2A-C maps to a conserved region, but does not affect a residue known to play a role in substrate binding or catalysis.

An alternative hypothesis is that the expression of heterologous PP2A-C subunit proteins down-regulates synthesis of endogenous C subunit protein. Loss of Pph21p expression would be tolerated in the presence of functional heterologous C subunits, but would be lethal in the presence of catalytically inactive mutants. A post-transcriptional autoregulatory mechanism limiting PP2A-C expression in mammalian cells has been reported recently (Baharians and Schonthal, 1998), and it is unclear whether a similar mechanism is operative in yeast. However, we have not observed down-regulation of endogenous C subunit protein levels in cultures expressing wild-type, dominant-defective, or non-complementing C subunits under *GAL* promoter control (D. Lizotte, A. DeLong, unpublished). Furthermore, a significant level of PP2A-C over-expression has been achieved in *S. pombe* (Kinoshita et al., 1993), and induction of high levels of PP2A-C mRNA is lethal in *S. cerevisiae*, suggesting that the C subunit protein is indeed over-expressed (Ronne et al., 1991). Thus down-regulation of endogenous C subunit levels does not account for the dominant-defective phenotype.

We have shown that yeast provides a functional assay

system for heterologous PP2A-C genes; this system is likely to be generalizable, given the high conservation of C subunit sequences. We have isolated dominant-defective mutants of PP2A genes from *Arabidopsis* and humans, and we plan to use these mutants to analyze PP2A functions in whole plants and in mammalian tissue culture cells. This genetic system may also prove useful for analyzing interactions between heterologous or homologous catalytic and regulatory subunits of PP2A.

### Acknowledgements

We thank S. Rundle, P. Philippsen, Z. Pitluk, N. Andjelkovic and B. Hemmings for plasmids, K. Arndt and B. Rockmill for yeast strains and advice on yeast genetics, and T. Stukenberg and M. Kirschner for the anti-PP2A antibody. We thank J. Sedivy for advice and comments on the manuscript, and M. Foulk, K. Yeung and A. Coleman for helpful discussions. We also thank K. Shamoun for assistance with site-directed mutagenesis. This work was supported by grants from the National Science Foundation (IBN-9604782), the NASA/NSF Joint program in Plant Biology (IBN-9416027) and the US Department of the Army (DAMD17-97-1-7113). The content of the information does not necessarily reflect the position or the policy of the government, and no official endorsement should be inferred. D.D.M.M. and H.R.C. were partially supported by Hughes/UTRA fellowships.

### References

- Ammerer, G., 1983. Expression of genes in yeast using the *ADCI* promoter. *Meth. Enzymol.* 101, 192–201.
- Ariño, J., Perez-Calderon, E., Cunillera, N., Camps, M., Posas, F., Ferrer, A., 1993. Protein phosphatases in higher plants: multiplicity of type 2A phosphatases in *Arabidopsis thaliana*. *Plant Mol. Biol.* 21 (3), 475–485.
- Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D.D., Seidman, J.G., Smith, J.A., Struhl, K., 1992. *Current Protocols in Molecular Biology*. Wiley-Interscience, New York.
- Baharians, Z., Schonthal, A.H., 1998. Autoregulation of protein phosphatase type 2A expression. *J. Biol. Chem.* 273, 19019–19024.
- Barford, D., 1996. Molecular mechanisms of the protein serine/threonine phosphatases. *Trends Biochem. Sci.* 21, 407–412.
- Cayla, X., Ballmer-Hofer, K., Merlevede, W., Goris, J., 1993. Phosphatase 2A associated with polyomavirus small-T or middle-T antigen is an okadaic acid-sensitive tyrosyl phosphatase. *Eur. J. Biochem.* 214, 281–286.
- Deng, W.P., Nickoloff, J., 1992. Site-directed mutagenesis of virtually any plasmid by elimination of a unique site. *Analyt. Biochem.* 200, 81–88.
- Evans, D.H.R., Stark, M.J.R., 1997. Mutations in the *Saccharomyces cerevisiae* type 2A protein phosphatase catalytic subunit reveal roles in cell wall integrity, actin cytoskeleton organization and mitosis. *Genetics* 145, 227–241.
- Garbers, C., DeLong, A., Deruère, J., Bernasconi and, P., Söll, D.,

1996. A mutation in protein phosphatase 2A regulatory subunit A affects auxin transport in *Arabidopsis*. *EMBO J.* 15, 2115–2124.
- Gietz, R.D., Sugino, A., 1988. New yeast–*Escherichia coli* shuttle vectors constructed with in vitro mutagenized yeast genes lacking six-base pair restriction sites. *Gene* 74, 527–534.
- Goldberg, J., Huang, H., Kwon, Y., Greengard, P., Nairn, A.C., Kuriyan, J., 1995. Three-dimensional structure of the catalytic subunit of protein serine/threonine phosphatase-1. *Nature* 376, 745–753.
- Götz, J., Probst, A., Ehler, E., Hemmings, B., Kues, W., 1998. Delayed embryonic lethality in mice lacking protein phosphatase 2A catalytic subunit  $\alpha$ . *Proc. Natl. Acad. Sci. USA* 95, 12370–12375.
- Healy, A.M., Zolnierowicz, S., Stapleton, A.E., Goebel, M., DePaoli-Roach, A.A., Pringle, J.R., 1991. *CDC55*, a *Saccharomyces cerevisiae* gene involved in cellular morphogenesis: identification, characterization and homology to the B subunit of mammalian type 2A protein phosphatase. *Mol. Cell. Biol.* 11, 5767–5780.
- Hemmings, B.A., Wernet, W., Mayer, R., Maurer, F., Hofsteenge, J., Stone, S.R., 1988. The nucleotide sequence of the cDNA encoding the human lung protein phosphatase 2A  $\beta$  catalytic subunit. *Nucleic Acids Res.* 16, 11366.
- Kaneko, S., Shima, H., Amagasa, T., Takagi, M., Sugimura, T., Nagao, M., 1995. Analysis by in vitro mutagenesis of PP2A  $\alpha$  okadaic acid responsive sequences. *Biochem. Biophys. Res. Commun.* 214, 518–523.
- Kinoshita, N., Ohkura, H., Yanagida, M., 1990. Distinct, essential roles of type 1 and 2A protein phosphatases in the control of the fission yeast cell division cycle. *Cell* 63, 405–415.
- Kinoshita, N., Yamano, H., Niwa, H., Yoshida, T., Yanagida, M., 1993. Negative regulation of mitosis by the fission yeast protein phosphatase ppa2. *Genes Dev.* 7, 1059–1071.
- Lec, T.H., Solomon, M.J., Mumby, M.C., Kirschner, M.W., 1991. INH, a negative regulator of MPF, is a form of protein phosphatase 2A. *Cell* 64, 415–423.
- Lin, F.C., Arndt, K.T., 1995. The role of *Saccharomyces cerevisiae* type 2A phosphatase in the actin cytoskeleton and in entry into mitosis. *EMBO J.* 14, 2745–2759.
- Mayer-Jackel, R.E., Hemmings, B.A., 1994. Protein phosphatase 2A, a 'ménage à trois'. *Trends Cell Biol.* 4, 287–291.
- Mayer-Jackel, R.E., Ohkura, H., Ferrigno, P., Andjelkovic, N., Shiomi, K., Uemura, T., Glover, D.M., Hemmings, B.A., 1994. *Drosophila* mutants in the 55 kDa regulatory subunit of protein phosphatase 2A show strongly reduced ability to dephosphorylate substrates of p34cdc2. *J. Cell Sci.* 107, 2609–2616.
- Mertz, P., Yuan, L., Sikkink, R., Rusnak, F., 1997. Kinetic and spectroscopic analysis of mutants of a conserved histidine in the metallophosphatases calcineurin and  $\lambda$  protein phosphatase. *J. Biol. Chem.* 272, 21296–21302.
- Mumby, M.C., Walter, G., 1993. Protein serine/threonine phosphatases: structure, regulation, and functions in cell growth. *Physiol. Rev.* 73 (4), 673–699.
- Murata, K., Wu, J., Brautigan, D.L., 1997. B cell receptor-associated protein  $\alpha 4$  displays rapamycin-sensitive binding directly to the catalytic subunit of protein phosphatase 2A. *Proc. Natl. Acad. Sci. USA* 94, 10624–10629.
- Pitluk, Z.W., McDonough, M., Sangan, P., Gonda, D.K., 1995. Novel CDC34 (UBC3) ubiquitin-conjugating enzyme mutants obtained by charge-to-alanine scanning mutagenesis. *Mol. Cell. Biol.* 15, 1210–1219.
- Ronne, H., Carlberg, M., Hu, G.Z., Nehlin, J.O., 1991. Protein phosphatase 2A in *Saccharomyces cerevisiae*: effects on cell growth and bud morphogenesis. *Mol. Cell. Biol.* 11, 4876–4884.
- Shima, H., Tohda, H., Aonuma, S., Nakayasu, M., DePaoli-Roach, A.A., Sugimura, T., Nagao, M., 1994. Characterization of the PP2A  $\alpha$  gene mutation in okadaic acid-resistant variants of CHO-K1 cells. *Proc. Natl. Acad. Sci. USA* 91, 9267–9271.
- Shu, Y., Yang, H., Hallberg, E., Hallberg, R., 1997. Molecular genetic analysis of Rts1p, a B' regulatory subunit of *Saccharomyces cerevisiae* protein phosphatase 2A. *Mol. Cell. Biol.* 17, 3242–3253.
- Smith, R.D., Walker, J.C., 1996. Plant protein phosphatases. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 47, 101–125.
- Snedden, A.A., Cohen, P.T.W., Stark, M.J.R., 1990. *S. cerevisiae* protein phosphatase 2A performs an essential cellular function and is encoded by two genes. *EMBO J.* 9, 4339–4346.
- Sontag, E., Fedorov, S., Kamibayashi, C., Robbins, D., Cobb, M., Mumby, M., 1993. The interaction of SV40 small tumor antigen with protein phosphatase 2A stimulates the map kinase pathway and induces cell proliferation. *Cell* 75, 887–897.
- Stamey, R.T., Rundle, S.J., 1996. Characterization of a novel isoform of a type 2A serine/threonine protein phosphatase from *Arabidopsis thaliana*. *Plant Physiol.* 110, 335.
- Stone, S.R., Mayer, R., Wernet, W., Maurer, F., Hofsteenge, J., Hemmings, B.A., 1988. The nucleotide sequence of the cDNA encoding the human lung protein phosphatase 2A  $\alpha$  catalytic subunit. *Nucleic Acids Res.* 16, 11365.
- van Zyl, W., Huang, W., Snedden, A.A., Stark, M., Camier, S., Werner, M., Marck, C., Sentenac, A., Broach, J.R., 1992. Inactivation of the protein phosphatase 2A regulatory subunit A results in morphological and transcriptional defects in *Saccharomyces cerevisiae*. *Mol. Cell Biol.* 12, 4946–4956.
- Wach, A., Brachat, A., Pohlmann, R., Philippsen, P., 1994. New heterologous modules for classical or PCR-based gene disruptions in *Saccharomyces cerevisiae*. *Yeast* 10, 1793–1808.
- Yang, S.I., Lickteig, R.L., Estes, R., Rundell, K., Walter, G., Mumby, M.C., 1991. Control of protein phosphatase 2A by simian virus 40 small-t antigen. *Mol. Cell. Biol.* 11, 1988–1995.
- Zhuo, S., Clemens, J.C., Stone, R.L., Dixon, J.E., 1994. Mutational analysis of a Ser/Thr phosphatase. Identification of residues important in phosphoesterase substrate binding and catalysis. *J. Biol. Chem.* 269, 26234–26238.

(contents continued)

|   |         |
|---|---------|
| <b>Linzmeier, R., Ho, C.H., Hoang, B.V. and Ganz, T.</b> .....  | 205-211 |
| A 450-kb contig of defensin genes on human chromosome 8p23  |         |
| <b>Huo, L. and Scarpulla, R.C.</b> .....  | 213-224 |
| Multiple 5'-untranslated exons in the nuclear respiratory factor 1 gene span 47 kb and contribute to transcript heterogeneity and translational efficiency  |         |
| <b>Sánchez, L.B., Morrison, H.G., Sogin, M.L. and Müller, M.</b> .....  | 225-231 |
| Cloning and sequencing of an acetyl-CoA synthetase (ADP-forming) gene from the amitochondriate protist, <i>Giardia lamblia</i>  |         |
| <b>Huang, W., Escribano, J., Sarfarazi, M. and Coca-Prados, M.</b> .....  | 233-240 |
| Identification, expression and chromosome localization of a human gene encoding a novel protein with similarity to the <i>pilB</i> family of transcriptional factors (pilin) and to bacterial peptide methionine sulfoxide reductases |         |
| <b>Miyazaki, S., Rasmussen, S., Imatani, A., Diella, F., Sullivan, D.T. and Callahan, R.</b> .....  | 241-247 |
| Characterization of the <i>Drosophila</i> ortholog of mouse <i>eIF-3p48/INT-6</i>   |         |
| <b>López, C.C., Kamnert, I., Scherbik, S.V. and Edström, J.-E.</b> .....  | 249-259 |
| Interspersed DNA element restricted to a specific group of telomeres in the dipteran <i>Chironomus pallidivittatus</i>  |         |
| <b>Avarbock, D., Salem, J., Li, L.-s., Wang, Z.-m. and Rubin, H.</b> .....  | 261-269 |
| Cloning and characterization of a bifunctional RelA/SpoT homologue from <i>Mycobacterium tuberculosis</i>   |         |
| <b>Sinha, M. and Hasan, G.</b> .....  | 271-276 |
| Sequencing and exon mapping of the <i>inositol 1,4,5-trisphosphate receptor</i> cDNA from <i>Drosophila</i> embryos suggests the presence of differentially regulated forms of RNA and protein  |         |
| <b>Author Index Vol. 233 (1999)</b> .....   | 277-278 |

**CONTENTS**  
**Direct**

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier Science books and journals. You can register for **ContentsDirect** online at: [www.elsevier.nl/locate/contentsdirect](http://www.elsevier.nl/locate/contentsdirect)